



International Journal of Orthopedics: Research & Therapy

Case Report

Atypical Presentation of Intra-Articular Osteoid Osteoma in the Proximal Femur: A Difficult Diagnosis - ②

James Choueiry and Jean-Emile Dubuc*

Department of Orthopedics, Saint-Luc University Hospital-UC Louvain, Avenue Hippocrate 10/2942, 1200 Brussels, Belgium

***Address for Correspondence:** Jean-Emile Dubuc, Department of Orthopedics, Saint-Luc University Hospital-UC Louvain, Avenue Hippocrate 10/2942, 1200 Brussels, Belgium, Tel: +32 (0)2 764 913; Fax: 32 (0)2 764 89 04; E-mail: jean-emile.dubuc@uclouvain.be

Submitted: 22 November 2017; **Approved:** 30 November 2017; **Published:** 01 December 2017

Cite this article: Choueiry J, Dubuc JE. Atypical Presentation of Intra-Articular Osteoid Osteoma in the Proximal Femur: A Difficult Diagnosis. Int J Ortho Res Ther. 2017;1(1): 025-027.

Copyright: © 2017 Choueiry J, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Osteoid osteoma is among the commonest bone tumors, primarily affecting young subjects. Often localized in the diaphysis cortex of long bones, the disease has a well-described symptomatology and imagery of choice for diagnosis. When in a different location, the diagnosis is less evident. We describe a case herein of an intra-articular osteoid osteoma of the hip misdiagnosed as a femoro-acetabular impingement and treated by means of hip arthroscopy.

INTRODUCTION

Osteoid Osteomas (OOs) are benign tumors consisting of a hypervascular inflammatory lytic nidus in the cortex of long bones. They are defined as benign Stage 2 lesions according to Musculoskeletal Society (MSTS) Staging. They often respond to aspirin, tending to disappear following several years of conservative management. The imagery of choice is Computed Tomography (CT), able to reveal the hypolucent nidus surrounded by an area of reactive sclerosis and periosteal thickening. Positron-Emission Tomography (PET) can reveal a double density sign, with the nidus the most intense surrounded by the less-intense bony sclerosis, giving the image of a halo. Magnetic Resonance Imaging (MRI) demonstrates the nidus with the surrounding bone edema. These findings may differ in cases involving atypical osteoma location.

CASE REPORT

A 19-year-old soccer player presented at consultation because of residual pain in the left hip 1 year after undergoing arthroscopy for femoroacetabular impingement. The patient had left inguinal nonirradiating pain, exacerbated by walking to the point of limping. Past pre-operative imagery revealed superior acetabular and posterior femoral-head osteophytes, suggesting pincer impingement and cam deformity. Post-operative imagery revealed excision of the superior osteophyte (Figure 1A, Figure 1B). In spite of this, the patient suffered from persistent pain, and a pelvic MRI was prescribed (Figure 2), revealing joint edema and synovial hypertrophy and an anterior cortical femoral neck lesion on STIR sequences. The lesion was hypolucent, surrounded by a hyperlucent rim. CT was performed for further evaluation, revealing a millimetric lesion that was slightly hyper intense and surrounded by a narrow osteolytic rim with no perinidal sclerotic bone reaction. Nevertheless, radiology was performed to confirm the diagnosis of an OO, with subsequent radiofrequency ablation. The patient was examined 5 months later and reported complete alleviation of pain and improvement in functional status. Physical examination showed a significantly improved range of motion for all movements except in internal rotation, where the hip remained stiff. Retrospectively, the lesion was not apparent on MRI prior to arthroscopy. On the other hand, the scan (Figure 3A) did reveal a subtle osteolytic lesion without the evident osteoid calcification that was present on the post-arthroscopy scan (Figure 3B). This could have been mistaken for an artefact.

DISCUSSION

Anatomically, OOs are typically located in the diaphysis cortex of long bones. In these locations, they present typical symptomatology and require normal imagery. In atypical locations, the diagnosis may become less evident. For example, patients with intraarticular hip lesions have less night pain [1], and imaging-based diagnosis proves more challenging. CT is not able to detect the sclerotic rim around the nidus, which has been attributed to absence of the bone-forming layer of the periosteum called the cambium layer [2]. The reactive sclerosis, however, may be a little distant [3]. X-ray can reveal




Figure 1: (A) Anteroposterior view of the pelvis with the hips in extension, revealing superior acetabular osteophytes and posterior femoral cam deformity on the left
(B) Excision of the acetabular osteophyte.




Figure 2: Axial cut of a T2-weighted image of the pelvis, revealing anterior cortical hypolucent lesion with concomitant bone and joint edema.




Figure 3: (A) Small intracortical lesion of the anterior border of the femoral neck shortly following manifestation of the patient's symptoms
(B) Increased size of the lesion with more osteoid calcification, yet no perinidal bone sclerosis.

distant reactive bony hypertrophy of the femoral neck, suggesting cam deformity. This can contribute to a patient's symptomatology, leading physicians away from the true cause. MRI can reveal labral tears, which are associated with cam deformity [4,5]. Bone edema can conceal the nidus, giving the appearance of regional osteoporosis or

an aggressive process [6,7]. The edema may extend to the synovium and soft tissues, mimicking inflammatory monarthritis or synovitis [8-11]. A small nidus can be masked by the intense signal exhibited by a bone edema on MRI [12], and a smaller MRI field is sometimes necessary, including half-moon sign assessment, as suggested by Klontas *et al.* [12,13]. Moreover, bone scintigraphy may not have diagnostic value in intra-articular lesions [14], and when lesions affect the feet or hands, they may be misdiagnosed as complex regional reflex dystrophy [15]. Symptoms can be acutely aggravated when OO contents perforate into the joint following cartilage erosion [16,17]. Briefly, OOs can occur anywhere in the bone, typically mimicking the usual pathologies of their surrounding area.

Children can also be affected, and can exhibit such an intense reactive bony sclerosis that this may obscure the site of the hypolucent nidus. In these cases, the CT scan slices need to be thinner than 2-3mm or the lesion may be missed [2,18]. With lesions located near the epiphysis, patients may present with limb-length discrepancy and axial deformation [14,19].

Sometimes, OOs may be mistaken for Brodie's abscesses. Osteoid nidus calcification, if present, looks the same as a sequestrum on imaging. On gadolinium-enhanced MRI, however, the nidus can be revealed as hypervascular and hyperintense, and the abscess may become a fistula, showing a break in the rim [20].

OOs may also be present in cancellous bone, the lesion potentially not associated with any reactive bone sclerosis process. CT scanning in these cases do not show the perinidal sclerotic contour. In fact, histologically, OOs can be intracortical (mainly), subperiosteal (e.g. talus) or cancellous. When cancellous, CT only reveals a small area of decreased bone density, while scintigraphy may not be able to detect the perinidal halo of a reactive bony sclerosis [14]. In contrast, dynamic contrast-enhanced MRI can reveal the early peak and washout flow of a hypervascular nidus, proving different from that of the surrounding bone edema, and may thus offer an alternative option in detecting atypical OOs [21,22].

Nevertheless, OOs remain benign tumors known to naturally regress over several years, with the use of medical management reported in several studies [23,24]. The regression may be caused by clotting of the vascularity inside the nidus [25]. In healed OOs, the sclerosis and cortical thickening may not disappear for many years, thus creating the potential for error [26,27]. Furthermore, the morphological appearance of femoroacetabular impingements does not necessarily mean they are the cause of hip pain, and careful assessment is thus essential [28].

REFERENCES

1. Radcliffe SN, Walsh HJ, Carty H. Osteoid osteoma: the difficult diagnosis. *Eur J Radiol.* 1998; 28: 67-79. <https://goo.gl/yUnVWA>
2. Ciftedemir M, Tuncel SA, Usta U. Atypical osteoid osteomas. *Eur J Orthop Surg Traumatol.* 2015; 25: 17-27. <https://goo.gl/6mCWby>
3. Cassar-Pullicino VN, McCall IW, Wan S. Intra-articular osteoid osteoma. *Clin Radiol* 1992; 45: 153-60. <https://goo.gl/Ciopa5>
4. Banga K, Racano A, Ayeni OR, Deheshi B. Atypical hip pain: coexistence of femoroacetabular impingement (FAI) and osteoid osteoma. *Knee Surg Sports Traumatol Arthrosc.* 2015; 23: 1571-4. <https://goo.gl/VKgMtv>
5. Ly JA, Coleman EM, Cohen GS, Kropf EJ. Unrecognized osteoid osteoma of the proximal femur with associated cam impingement. *J Hip Preserv Surg.* 2016; 3: 236-7. <https://goo.gl/HZDqZ9>
6. Davies M, Cassar-Pullicino VN, Davies AM, McCall IW, Tyrrell PN. The diagnostic accuracy of MR imaging in osteoid osteoma. *Skeletal Radiol.* 2002; 31: 559-69. <https://goo.gl/x8fLRB>

7. Wiener SN, Kirschenbaum D. Osteoid osteoma presenting as regional osteoporosis. *Clin Nucl Med.* 1980; 5: 68-9. <https://goo.gl/gb3nuu>
8. Snarr JW, Abell MR, Martel W. Lymphofollicular synovitis with osteoid osteoma. *Radiology.* 1973; 106: 557-60. <https://goo.gl/3NLyKN>
9. Barbiera F, Bartolotta TV, Lo Casto A, Pardo S, Rossello M, De Maria M. Intra-articular osteoid osteoma: diagnostic imaging in three cases. *Radiol Med.* 2002; 103: 464-73. <https://goo.gl/2oA8FM>
10. Herget GW, Sudkamp NP, Bohm J, Helwig P. Osteoid osteoma of the femoral neck mimicking monarthritis and causing femoroacetabular impingement. *Acta Chir Orthop Traumatol Cech.* 2012; 79: 275-8. <https://goo.gl/JH7S2x>
11. Thompson GH, Wong KM, Konsens RM, Vibhakar S. Magnetic resonance imaging of an osteoid osteoma of the proximal femur: a potentially confusing appearance. *J Pediatr Orthop.* 1990; 10: 800-4. <https://goo.gl/8q1Bgh>
12. James SL, Panicek DM, Davies AM. Bone marrow oedema associated with benign and malignant bone tumours. *Eur J Radiol.* 2008; 67: 11-21. <https://goo.gl/mcFZEF>
13. Klontzas ME, Zibis AH, Karantanas AH. Osteoid Osteoma of the Femoral Neck: Use of the Half-Moon Sign in MRI Diagnosis. *AJR Am J Roentgenol.* 2015; 205: 353-7. <https://goo.gl/ivX39F>
14. Szendroi M, Kollo K, Antal I, Lakatos J, Szoke G. Intraarticular osteoid osteoma: clinical features, imaging results, and comparison with extraarticular localization. *J Rheumatol.* 2004; 31: 957-64. <https://goo.gl/zoxTTD>
15. Cakar E, Durmus O, Kiralp MZ, Dincer U. An unusual case of osteoid osteoma misdiagnosed as inflammatory joint disease and complex regional pain syndrome I. *Acta Reumatol Port.* 2009; 34: 670-1. <https://goo.gl/VSRgJ4>
16. Bhardwaj P, Sharma C, Sabapathy SR. Synovitis of the wrist joint caused by an intraarticular perforation of an osteoid osteoma of the scaphoid. *Indian J Orthop.* 2012; 46: 599-601. <https://goo.gl/2weXP4>
17. De Smet L. Synovitis of the wrist joint caused by an intraarticular perforation of an osteoid osteoma of the radial styloid. *Clin Rheumatol.* 2000; 19: 229-30. <https://goo.gl/HSVGjU>
18. Schlur C, Bachy M, Wajfisz A, Ducou le Pointe H, Josset P, Vialle R. Osteoid osteoma mimicking Brodie's abscess in a 13-year-old girl. *Pediatr Int.* 2013; 55: 29-31. <https://goo.gl/W4atj8>
19. Gabriel H, Fitzgerald SW, Myers MT, Donaldson JS, Poznanski AK. MR imaging of hip disorders. *Radiographics.* 1994; 14: 763-81. <https://goo.gl/wWCNUK>
20. McGrath BE, Bush CH, Nelson TE, Scarborough MT. Evaluation of suspected osteoid osteoma. *Clin Orthop Relat Res.* 1996; 327: 247-52. <https://goo.gl/yHuVd1>
21. Liu PT, Chivers FS, Roberts CC, Schultz CJ, Beauchamp CP. Imaging of osteoid osteoma with dynamic gadolinium-enhanced MR imaging. *Radiology.* 2003; 227: 691-700. <https://goo.gl/RcWGV8>
22. Zampa V, Bargellini I, Ortori S, Faggioni L, Cioni R, Bartolozzi C. Osteoid osteoma in atypical locations: the added value of dynamic gadolinium-enhanced MR imaging. *Eur J Radiol.* 2009; 71: 527-35. <https://goo.gl/aQCwW>
23. Kneisl JS, Simon MA. Medical management compared with operative treatment for osteoid-osteoma. *J Bone Joint Surg Am.* 1992; 74: 179-85. <https://goo.gl/fzZaQk>
24. Saville PD. A medical option for the treatment of osteoid osteoma. *Arthritis Rheum.* 1980; 23: 1409-11. <https://goo.gl/jY5Rgg>
25. Papanthassiou ZG, Megas P, Petsas T, Papachristou DJ, Nilas J, Siablis D. Osteoid osteoma: diagnosis and treatment. *Orthopedics.* 2008; 31: 1118. <https://goo.gl/HkMb9x>
26. Freiburger RH, Loitman BS, Helpert M, Thompson TC. Osteoid osteoma; a report on 80 cases. *Am J Roentgenol Radium Ther Nucl Med.* 1959; 82: 194-205. <https://goo.gl/5iGMmv>
27. Vickers CW, Pugh DC, Ivins JC. Osteoid osteoma; a fifteen-year follow-up of an untreated patient. *J Bone Joint Surg Am.* 1959; 41: 357-8. <https://goo.gl/BqHHUA>
28. Mainzer J, Ganz R, Tibor L, Leunig M. Not all hip pain is impingement: femoral neck osteoid osteoma in a patient with a coexisting cam deformity. *JBJS Case Connect.* 2012; 2: e311-5. <https://goo.gl/Xfnci>